

VALUES OF ZETA FUNCTIONS AT NEGATIVE INTEGERS, DEDEKIND SUMS AND TORIC GEOMETRY

STAVROS GAROUFALIDIS AND JAMES E. POMMERSHEIM

Dedicated to our teacher, W. Fulton.

ABSTRACT. We study relations among special values of zeta functions, invariants of toric varieties, and generalized Dedekind sums. In particular, we use invariants arising in the Todd class of a toric variety to give a new explicit formula for the values of the zeta function of a real quadratic field at nonpositive integers. We also express these invariants in terms of the generalized Dedekind sums studied previously by several authors. The paper includes conceptual proofs of these relations and explicit computations of the various zeta values and Dedekind sums involved.

CONTENTS

1. Introduction	2
1.1. Zeta functions	2
1.2. Toric geometry	4
1.3. Dedekind sums	6
1.4. Is toric geometry needed?	7
1.5. Plan of the proof	8
1.6. Acknowledgments	8
2. The Todd power series of a cone	8
2.1. General properties of the Todd power series	9
2.2. Properties of Todd power series of two-dimensional cones	11
2.3. Continued fraction expansion for the Todd power series	12
2.4. Zeta function values in terms of Todd power series	14
3. Zeta functions of number fields	16
3.1. Parametrizing triples (M_n, Q_b, τ_b)	16
3.2. Proof of Theorem 1	19
4. Dedekind sums in terms of Todd power series	21
References	22

Date: This edition: September 12, 1998; First edition: November 22, 1997 .

The authors were partially supported by NSF grants DMS-95-05105 and DMS-95-08972, respectively. This and related preprints can also be obtained at <http://www.math.brown.edu/~stavrosg> 1991 *Mathematics Classification*. Primary 11M06. Secondary 14M25, 11F20.

Key words and phrases: zeta functions, Dedekind sums, toric varieties.

1. INTRODUCTION

In the present paper, we study relations among special values of zeta functions of real quadratic fields, properties of generalized Dedekind sums and Todd classes of toric varieties. The main theme of the paper is the use of toric geometry to explain in a conceptual way properties of the values of zeta functions and Dedekind sums, as well to provide explicit computations.

Both toric varieties and zeta functions associate numerical invariants to cones in lattices; however, with different motivations and applications. Though we will focus on the case of two-dimensional cones in the present paper, we introduce notation and definitions that are valid for cones of arbitrary dimension. The reasons for this added generality is clarity, as well as preparation for the results of a subsequent publication.

1.1. Zeta functions. We begin by reviewing the first source of numerical invariants of cones: the study of zeta functions.

Given a number field K , its zeta function is defined (for $\operatorname{Re}(s)$ sufficiently large) by:

$$\zeta(K, s) = \sum_{\alpha} \frac{1}{Q(\alpha)^s}$$

where the summation is over all nonzero ideals, and Q is the norm. The above function admits a meromorphic continuation in $\mathbb{C}$, with a simple pole at $s = 1$, and regular everywhere else. Lichtenbaum [Li] conjectured a specific behavior of the zeta function $\zeta(K, s)$ at nonpositive integers related to the global arithmetic of the number field. In the special case of a totally real number field K , Lichtenbaum conjectured that the values of the zeta function at negative integers are rational numbers which involve the rank of the algebraic (or étale) K -theory of K . It turns out (see [Sh1] and [Za2, Section 2]) that for a totally real field K the zeta function can be decomposed as a sum $\zeta(K, s) = \sum_I \zeta(I, s)$ where the sum is over the (finite set of) narrow ideal classes. For each narrow ideal class I , there is a finite set $\{\sigma_{i,I}^\circ\}_i$ of open cones in a rank $[K : \mathbb{Q}]$ lattice M_I in K such that $\zeta(I^{-1}, s) = \sum_i \zeta_{Q, \sigma_{i,I}^\circ}(s)$ where Q is a multiple of the norm on K and where for an open cone τ° we have

$$\zeta_{Q, \tau^\circ}(s) = \sum_{a \in \tau^\circ \cap M_I} \frac{1}{Q(a)^s}.$$

The problem of calculating the zeta values $\zeta_{Q, \tau^\circ}(-n)$ for $n \geq 0$ for all triples (M, Q, τ°) that arise from totally real fields has attracted a lot of attention by several authors. Klingen [Kl] and Siegel [Si1, Si2] (using analytic methods) proved that the values of the zeta functions of totally real fields at nonpositive integers are rational numbers and provided an algorithm for calculating them. Meanwhile, Shintani [Sh1] (using algebraic and combinatorial methods) gave an independent calculation of the zeta values $\zeta_{Q, \tau^\circ}(-n)$ for an arbitrary Q which is a product of linear forms. Meyer [Me] and Zagier [Za3] gave another calculation of the zeta values $\zeta_{Q, \tau^\circ}(-n)$ for rank two lattices. Related results have also been obtained by P. Cassou-Noguès, [C-N1, C-N2, C-N3]. More recently, Hayes [Hs], Sczech [Sc1, Sc2] and Stevens [St] have constructed $\operatorname{PGL}_m(\mathbb{Q})$ cocycles which, among other things, provide a calculation of the zeta values (at nonpositive integers) of totally real number fields in terms of generalized Dedekind sums.

We now specialize to the case of real quadratic fields. Given a narrow ideal class I , there is a cone σ_I (i.e., the convex hull of two rays from the origin) in a rank two lattice $M \subseteq K$,

such that: $\zeta(I^{-1}, s) = \zeta_{Q, \sigma_I}(s)$. Here Q is a multiple of the norm form and $\zeta_{Q, \sigma}$ is given by the following definition:

Definition 1.1. Let M be an two-dimensional lattice, and let V denote the real vector space $M_{\mathbb{R}} = M \otimes \mathbb{R}$. Let $Q : M_{\mathbb{R}} \rightarrow \mathbb{R}$ be a *nonzero quadratic homogenous function* i.e., a function satisfying $Q(av) = a^2Q(v)$ for $a \in \mathbb{R}, v \in M_{\mathbb{R}}$ and such that Q is not identically 0. We set

$$(1) \quad \zeta_{Q, \tau}(s) = \sum_{a \in \tau \cap M} \frac{wt(\tau, a)}{Q(a)^s}$$

where τ is a rational two-dimensional cone in M and $wt(\tau, \cdot) : M \rightarrow \mathbb{Q}$ is the weight function defined by:

$$(2) \quad wt(\tau, a) = \begin{cases} 1 & \text{if } a \text{ lies in the interior of } \tau \\ 1/2 & \text{if } a \text{ lies in the boundary of } \tau, \text{ and } a \neq 0 \\ 0 & \text{otherwise.} \end{cases}$$

We will assume throughout that Q is positive on all of τ ; that is, for all $a \in \tau$, $a \neq 0$, we have $Q(a) > 0$. The above zeta function, defined for $Re(s)$ sufficiently large, [Za4] can be analytically continued to a meromorphic function on $\mathbb{C}$, regular everywhere except at 1.

In [Za4], Zagier showed that after possibly multiplying M by a totally positive number and Q by a nonzero rational number (which only multiplies values of the zeta function by a nonzero constant), every triple (M, Q, τ) which arise from real quadratic field can be constructed by means of a finite sequence $b = (b_0, \dots, b_{r-1})$ of integers greater than 1 and not all equal to 2, which will be denoted by (M_b, Q_b, τ_b) . In addition, there are canonical vectors $A_0, \dots, A_r$ in M_b (that depend on b) that subdivide the cone $\tau_b = \langle A_0, A_r \rangle$ (i.e., the cone whose extreme rays are A_0 and A_r) in M_b in r nonsingular cones $\langle A_i, A_{i+1} \rangle$. For a detailed discussion, see Section 3.1.

We will now give an explicit formula for the zeta values $\zeta_{Q_b, \tau_b}(-n)$ (for $n \geq 0$), which in turn enables one to calculate the values of the zeta function of a real quadratic field at nonpositive integers. While both the motivation and the proof involve concepts from the theory of toric geometry, the formula can be stated and understood without a knowledge of toric varieties. We will state the formula here, and in the next subsection, we will discuss concepts from toric geometry which are necessary for the proof and which lead to a conceptual understanding of the present formula.

Let λ_m be defined by the power series:

$$(3) \quad \frac{h}{1 - e^{-h}} = \sum_{m=0}^{\infty} \lambda_m h^m$$

thus we have: $\lambda_m = (-1)^m B_m / m!$ where B_m is the m^{th} Bernoulli number. (See also Definition 1.4 below.) Note that if $m > 1$ is odd, then $\lambda_m = 0$. For $d \geq 2$ even, define homogeneous polynomials $P_d(X, Y), R_d(X, Y)$ of degree $d - 2$ by:

$$\begin{aligned} P_d(X, Y) &= \epsilon_d \sum_{i+j=d, i, j > 0} \lambda_i \lambda_j X^{i-1} Y^{j-1}, \\ R_d(X, Y) &= \frac{X^{d-1} + Y^{d-1}}{X + Y} = X^{d-2} - X^{d-3}Y + \dots + Y^{d-2}, \end{aligned}$$

where $\epsilon_2 = -1$ and $\epsilon_d = 1$ for all even $d > 2$.

We then have:

Theorem 1. *The values $\zeta_{Q_b, \tau_b}(-n)$ for $n \geq 0$ are polynomials in b_i with rational coefficients (symmetric under cyclic permutation of the b_i) given explicitly as follows:*

$$(4) \quad \zeta_{Q_b, \tau_b}(-n) = (-1)^n n! \left\{ \sum_{i=1}^r P_{2n+2} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) \diamond e^{-Q_b(xA_{i-1} + yA_i)} \right. \\ (5) \quad \left. + \lambda_{2n+2} \sum_{i=1}^r b_i R_{2n+2} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) \diamond e^{-Q_b(xA_{i-1} + yA_{i+1})} \right\}$$

The diamond symbol indicates the evaluation of a partial derivative at the origin.

In particular, we obtain the formula of Zagier [Za1, Equation 3.3]:

$$(6) \quad \zeta_{Q_b, \tau_b}(0) = \frac{1}{12} \sum_{i=0}^{r-1} (b_i - 3).$$

1.2. Toric geometry. The second source of numerical invariants of cones comes from the theory of *toric varieties*. For a general reference on toric varieties, see [Da] or [Fu].

Founded in the 1970s, the subject of toric varieties provides a strong link between algebraic geometry and the theory of convex bodies in a lattice. To each lattice polytope (the convex hull of a finite set of lattice points) is associated an algebraic variety with a natural torus action. This correspondence enables one to translate important properties and theorems about lattice polytopes into the language of algebraic geometry, and vice-versa. One important example of this is the very classical problem of counting the number of lattice points in a polytope. The early pioneers in the subject of toric varieties found that this problem, translated into algebraic geometry, becomes the problem of finding the Todd class of a toric variety. Much progress has been made over the past ten years on the Todd class problem, and this has led to a greater understanding of the lattice point counting question.

One approach to computing the Todd class of a toric variety is the fundamental work of R. Morelli [Mo]. He settled a question of Danilov by proving a local formula expressing the Todd class of a toric variety as a cycle. Another approach, introduced in by the first author in [P1, P2], is to express the Todd class as a polynomial in the torus-invariant cycles. Dedekind sums appear as coefficients in these polynomials, and this leads to lattice point formulas in terms of Dedekind sums, as well as new reciprocity laws for Dedekind sums [P1]. Cappell and Shaneson [CS] subsequently announced an extension of the program of [P1] in which they proposed formulas for the Todd class of a toric variety in all dimensions. In [P3] it was shown that the polynomials of [P1, P2] can be expressed nicely as the truncation of a certain power series whose coefficients were shown to be polynomial-time computable using an idea of Barvinok [Ba]. A beautiful power series expression for the equivariant Todd class of a toric variety was given by Brion and Vergne in [BV2]. Guillemin [Gu] also proved similar Todd class formulas from a symplectic geometry point of view. Furthermore, in [BV1], Brion and Vergne use the Todd power series of [BV2] to give a formula for summing any polynomial function over a polytope.

The power series studied in [P3] and [BV1, BV2] are, in fact, identical (see Section 2) and play a central role in the present paper. A detailed discussion of the properties of these

power series, which we call the *Todd power series of a cone*, is contained in Section 2. In this section, we state two theorems about the Todd power series of a two-dimensional cone that we will need in our study of the zeta function.

Given independent rays $\rho_1, \dots, \rho_n$ from the origin in an n -dimensional lattice N , the convex hull of the rays in the vector space $N_{\mathbb{R}} = N \otimes \mathbb{R}$ forms an n -dimensional cone $\sigma = \langle \rho_1, \dots, \rho_n \rangle$. Cones of this type (that is, ones generated by linearly independent rays) are called *simplicial*. Let $\mathcal{C}^n(N)$ denote the set of n -dimensional simplicial cones of N with ordered rays. There is then a canonical function

$$\mathbf{t} : \mathcal{C}^n(N) \rightarrow \mathbb{Q}[[x_1, \dots, x_n]],$$

invariant under lattice automorphisms, which associates to each cone σ a power series $\mathbf{t}_{\sigma}$ with rational coefficients, called the Todd power series of σ . Several ways of characterizing this function are given in Section 2. These include an N -additivity property (See Proposition 2.1), an exponential sum over the cone (Proposition 2.3) and an explicit cyclotomic sum formula (Proposition 2.4.)

In the case of a two-dimensional cone σ , the coefficient of xy of $\mathbf{t}_{\sigma}(x, y)$ was identified as a *Dedekind sum* [P1]. Furthermore, in [P1] it was shown that *reciprocity formulas* for Dedekind sums follow from an N -additivity formula for $\mathbf{t}$. Zagier's higher-dimensional Dedekind sums [Za1] were later shown to appear as coefficients [BV2]. It is natural to conjecture that, in the two-dimensional case, *all* the coefficients of $\mathbf{t}$ are generalized Dedekind sums and that reciprocity properties of generalized Dedekind sums will be related to the N -additivity formula of $\mathbf{t}$. Indeed, this is the case: see Theorem 4.

We now present an explicit link between Todd power series and zeta functions. The following theorem expresses the values of zeta function at negative integers in terms of the Todd power series of a two-dimensional cone. Note that the idea of considering the Todd power series as a differential operator, and applying this to an integral over a shifted cone is not new: this was introduced in [KP] and developed further in [BV1].

First we introduce some notation which is standard in the theory of toric varieties. If τ is a cone in a lattice M , the dual cone $\check{\tau}$ is a cone in the dual lattice $N = \text{Hom}(M, \mathbb{Z})$, defined by

$$\check{\tau} = \{v \in N \mid \langle v, u \rangle \geq 0 \text{ for all } u \in \tau\}.$$

The dual of an n -dimensional simplicial cone $\tau = \langle \rho_1, \dots, \rho_n \rangle$ is generated by the rays u_i a normal to the $n - 1$ -dimensional faces of τ ; hence $\check{\tau}$ is also a simplicial cone. Given $h = (h_1, \dots, h_n) \in \mathbb{R}^n$, we define the *shifted cone* $\tau(h)$ to be the following cone in $M_{\mathbb{R}}$:

$$\tau(h) = \{m \in M_{\mathbb{R}} \mid \langle u_i, m \rangle \geq -h_i \text{ for all } i = 1, \dots, n\}.$$

Here (and throughout) we have identified each ray u_i with the primitive lattice point on that ray, that is, the nonzero lattice point on u_i closest to the origin.

Given an n -dimensional simplicial cone $\tau = \langle \rho_1, \dots, \rho_n \rangle$ in an n -dimensional lattice M , the *multiplicity* of τ , denoted by $\text{mult}(\tau)$, is defined to be the index in M of the sublattice $\mathbb{Z}\rho_1 + \dots + \mathbb{Z}\rho_n$ (again identifying the rays ρ_i with their primitive lattice points.) Thus the multiplicity of τ is simply the volume of the parallelepiped formed by the vectors from the origin to the primitive lattice points on the rays of τ .

Theorem 2. *Let τ be a two-dimensional cone of multiplicity q in a two-dimensional lattice M , and let $\sigma = \check{\tau}$ be the dual cone in $N = \text{Hom}(M, \mathbb{Z})$. Then for all $n \geq 0$, we have:*

$$\zeta_{Q,\tau}(-n) = (-1)^n n! \left\{ (\mathbf{t}_\sigma)_{(2n+2)} \left(\frac{\partial}{\partial h_1}, \frac{\partial}{\partial h_2} \right) - \delta_{n,0} \frac{q}{2} \frac{\partial^2}{\partial h_1 \partial h_2} \right\} \diamond \int_{\tau(h)} \exp(-Q(u)) du,$$

where all derivatives above are evaluated at $h_1 = h_2 = 0$ and $\delta_{n,0} = 1$ (resp. 0) if $n = 0$ (resp. $n \neq 0$).

In this equation, $(\mathbf{t}_\sigma)_d$ denotes the degree d part of the Todd power series thought of as an (infinite order) constant coefficients differential operator acting on the function $h \rightarrow \int_{\tau(h)} \exp(-Q(u)) du$.

The coefficients of $\mathbf{t}_\sigma$ for a two-dimensional cone may be expressed explicitly in terms of continued fractions. We now state this formula.

Let σ be a two-dimensional cone in a lattice N . Then there exist a (unique) pair of relatively prime integers p, q with $0 < p \leq q$ such that σ is lattice equivalent (equivalent under the automorphism group of the lattice) to the cone $\sigma_{(p,q)} \stackrel{\text{def}}{=} \langle (1, 0), (p, q) \rangle$ in $\mathbb{Z}^2$. Such a cone will be called a cone of type (p, q) . Let b_i, h_i, k_i, X_i are defined in terms of the negative-regular continued fraction expansions:

$$(7) \quad \frac{q}{p} = [b_1, \dots, b_{r-1}], \quad \frac{h_i}{k_i} \stackrel{\text{def}}{=} [b_1, \dots, b_{i-1}], \quad X_i \stackrel{\text{def}}{=} -h_i x + (qk_i - ph_i)y.$$

(Throughout, we use such bracketed lists to denote finite negative-regular continued fractions.) We then have the following continued fraction expression for the degree d part $(\mathbf{t}_\sigma)_d$ of the Todd power series $\mathbf{t}_\sigma$.

Theorem 3. *For σ of type (p, q) as above, and for $d \geq 2$ an even integer, we have:*

$$\begin{aligned} (\mathbf{t}_\sigma)_d(x, y) &= qxy \sum_{i=1}^r P_d(X_{i-1}, X_i) + \lambda_d qxy \sum_{i=1}^{r-1} b_i R_d(X_{i-1}, X_{i+1}) \\ &\quad - \lambda_d (xX_1^{d-1} + yX_{r-1}^{d-1}) + \frac{1}{2} \delta_{d,2} qxy. \end{aligned}$$

If $d \geq 1$ is odd, then $(\mathbf{t}_\sigma)_d(x, y) = \frac{1}{2} \lambda_{d-1} q^{d-2} xy (x^{d-2} + y^{d-2})$.

Remark 1.2. We have stated the formula above in the form we will need it for our study of zeta functions. However, from the toric geometry point of view, it is more natural to use the continued fraction expansion of $\frac{q}{q-p}$ instead. This formula will be given in Section 2.3.

Remark 1.3. The formula for the Todd operator in Theorem 3 is reminiscent of formulas in quantum cohomology, even though we know of no conceptual explanation for this fact.

1.3. Dedekind sums. Finally, we calculate the coefficients of the Todd power series of two-dimensional cones in terms of a particular generalization $s_{i,j}$ (defined below) of the classical Dedekind sums. For an excellent review of the properties of the classical Dedekind sum, see [R-G]. Several generalizations of the classical Dedekind sums were studied by T. Apostol, Carlitz, C. Meyer, D. Solomon, and more recently and generally by U. Halbritter [AV, Ha, Me, So]. These papers investigate the sums (and several other generalizations of them, which we will not consider here) given in the following definition:

Definition 1.4. For relatively prime integers p, q (with $q \neq 0$), and for nonnegative integers i, j , we define the following *generalized Dedekind sum*:

$$(8) \quad s_{i,j}(p, q) = \frac{1}{i!j!} \sum_{a=1}^q \widehat{\text{Ber}}_i\left(\left\langle \frac{a}{q} \right\rangle\right) \widehat{\text{Ber}}_j\left(\left\langle -\frac{ap}{q} \right\rangle\right)$$

where for a real number x , we denote by $\langle x \rangle$ the (unique) number such that $\langle x \rangle \in (x + \mathbb{Z}) \cap (0, 1]$. Here Ber_m denotes the m^{th} Bernoulli polynomial, defined by the power series $\sum_{m=0}^{\infty} \text{Ber}_m(x) \frac{t^m}{m!} = \frac{te^{tx}}{1-e^t}$, and $\widehat{\text{Ber}}_m$ denotes the restriction of the m^{th} Bernoulli polynomial Ber_m to $(0, 1]$, with the boundary condition $\widehat{\text{Ber}}_m(1) \stackrel{\text{def}}{=} \frac{1}{2}(\text{Ber}_m(1) + \text{Ber}_m(0)) = B_m + \delta_{m,1}/2$, where B_m is the m^{th} Bernoulli number, defined by $B_m = \text{Ber}_m(0)$.¹

By definition, the sum $s_{1,1}(-p, q)$ coincides with the classical Dedekind sum $s(p, q)$ (cf. [R-G].) An important property of generalized Dedekind sums is a reciprocity formula which leads to an evaluation in terms of negative-regular continued fractions. This reciprocity formula was most conveniently written by D. Solomon in terms of an additivity formula of a generating power series [So, Theorem 3.3]. On the other hand, the Todd operator also satisfies an additivity property. We denote by $f_{i,j}(p, q)$ (for nonnegative integers i, j) the coefficient of $x^i y^j$ of the power series $\mathbf{t}_{\sigma(p,q)}(x, y)$ (abbreviated by $\mathbf{t}_{(p,q)}(x, y)$). We then have:

Theorem 4. Let $p, q \in \mathbb{Z}$ be relatively prime with $q \neq 0$. If $i, j > 1$, then we have:

$$(9) \quad f_{i,j}(p, q) = q^{i+j-1} (-1)^i s_{i,j}(p, q).$$

If $i = 1$ or $j = 1$, the above equation is true when the correction term

$$q^{i+j-1} (-1)^{i+j} \frac{B_i B_j}{i!j!}$$

is added to the right hand side.

Corollary 1.5. Fixing r , for all nonnegative integers i, j and with the notation of (7), $s_{i,j}(p, q)$ are polynomials in $b_i, 1/q$.

Remark 1.6. For $i = j = 1$, the above theorem was obtained in [P1], and was a motivation for the results of the present paper.

1.4. Is toric geometry needed? Some natural questions arise, at this point:

- Is toric geometry needed?
- How do the statement and proof of Theorem 1 differ from the statement and proof of Zagier's formula [Za4]?

With respect to the first question, Theorem 1 (an evaluation of zeta functions associated to real quadratic fields) is stated without reference to toric geometry, and a close examination reveals that its proof is based on an analytic Lemma 2.11 and on the two dimensional analogue of the Euler-MacLaurin formula given by Proposition 2.13. In addition, two dimensional cones can be canonically subdivided into nonsingular cones, so in a sense the two

¹There seem to be two conventions for denoting Bernoulli numbers; one, which we follow here, is used often in number theory texts. Due to the facts that $B_{2n+1} = 0$ for $n \geq 1$, and that the sign of B_{2n} alternates, the other convention, which is often used in intersection theory, defines the n^{th} Bernoulli number to be $(-1)^{n+1} B_{2n}$.

dimensional analogue of the Euler-MacLaurin formula can be obtained by the classical (one dimensional) one, as is used by Zagier [Za4]. Furthermore, number theory offers, for every $m \geq 2$, a canonical *Eisenstein cocycle* of $\mathrm{PGL}_m(\mathbb{Q})$ [Sc1, Sc2] that expresses, among other things, the generalized Dedekind sums $s_{i,j}$ in terms of negative-regular continued fraction expansions like the ones of Theorems 3 and 4, and the values (at nonpositive integers) of zeta functions of totally real fields (of degree m) in terms of generalized Dedekind sums. See also [So, St].

On the other hand, toric geometry constructs for every simplicial cone σ in an m -dimensional lattice N , a canonical Todd power series $\mathbf{t}_\sigma$ satisfying an N -additivity property (see Proposition 2.1 below). The coefficients of the Todd power series are generalized Dedekind sums, and the power series itself is intimately related to the Euler-MacLaurin summation formula. As a result, for $m = 2$, we provide a toric geometry explanation of Theorems 1 and 4.

With respect to the second question, Zagier [Za4] obtained similar formulas for the values of the zeta function of a quadratic number field at negative integers. Zagier used additivity in the lattice M , whereas we use additivity in the dual lattice N . N -additivity implies that these zeta values are a priori polynomials in b_i and $1/q$; moreover these polynomials are directly defined in terms of the Todd operator and the quadratic form Q . Thus the present paper is an example of the rather subtle difference between N -additivity and M -additivity. This difference can be seen by contrasting the additivity property of the zeta function $\zeta_{Q,\tau}$ with the additivity property satisfied by Todd operator. The modified Todd operator $\mathbf{s}_\sigma$ (introduced in Section 2.1) is N -additive under subdivisions of the cone σ , i.e., it satisfies equation (11) below. On the other hand, the function $\zeta_{Q,\cdot}(s) : \mathcal{C}^2(M) \rightarrow \mathbb{C}$ is M -additive, i.e., it satisfies:

$$(10) \quad \zeta_{Q,\tau}(s) = \zeta_{Q,\tau_1}(s) + \zeta_{Q,\tau_2}(s)$$

whenever τ is subdivided into cones τ_1 and τ_2 . This M -additivity of $\zeta_{Q,\cdot}(s)$ is due to the particular choice of the weight function wt , and is purely set-theoretic. In contrast, the N -additivity of $\mathbf{s}_\sigma$ is somewhat deeper, and has the advantage that only cones of maximal dimension are involved.

1.5. Plan of the proof. In Section 2, we review well known properties of Todd power series and prove Theorems 2 and 3. In Section 3, we review the relation between the zeta functions that we consider and the zeta functions of real quadratic fields. We give a detailed construction of zeta functions, and prove Theorem 1. Finally, in Section 4 we review properties of generalized Dedekind sums and prove Theorem 4.

1.6. Acknowledgments. We wish to thank M. Rosen for encouraging conversations during the academic year 1995-96. We also thank B. Sturmfels and M. Brion for their guidance. We especially wish to thank W. Fulton for enlightening and encouraging conversations since our early years of graduate studies.

2. THE TODD POWER SERIES OF A CONE

In this section we study properties of the Todd power series $\mathbf{t}_\sigma$ associated to a simplicial cone σ . Section 2.1 provides an introduction and statements of several previously discovered formulas for the Todd power series. In Section 2.2 we make these formulas explicit for two-dimensional cones. Section 2.3 contains a proof of the explicit continued fraction formula for the Todd power series of a two-dimensional cone. Finally, in Section 2.4 we prove Theorem 2

which links Todd power series with the problem of evaluating zeta functions on nonpositive integers.

2.1. General properties of the Todd power series. Todd power series were studied in connection with the Todd class of a simplicial toric variety in [P3]. Independently, they were introduced in [BV2] in the study of the equivariant Todd class of a simplicial toric variety. In addition, these power series appear in Brion and Vergne's formula for counting lattice points in a simple polytope [BV1], which is an extension of Khovanskii and Pukhlikov's formula [KP]. In these remarkable formulas, the power series in question are considered as differential operators which are applied to the volume of a deformed polytope. The result yields the number of lattice points in the polytope, or more generally, the sum of any polynomial function over the lattice points in the polytope. Below (Proposition 2.13), we give a version of this formula expressing the sum of certain functions over the lattice points contained in a simplicial cone.

The Todd power series considered in the works cited above are also closely related to the fundamental work of R. Morelli on the Todd class of a toric variety. [Mo]. A precise connection is given in [P3, Section 1.8].

We now state some of the properties of the Todd power series of a simplicial cone. Our purpose is twofold: we will need these properties in our application to zeta functions, and we wish to unite the approaches of the works cited above. Here, we follow the notation of [P3].

Let $\sigma = \langle \rho_1, \dots, \rho_n \rangle$ be an n -dimensional simplicial cone in an n -dimensional lattice N . The Todd power series $\mathbf{t}_\sigma$ of σ is a power series with rational coefficients in variables $x_1, \dots, x_n$ corresponding to the rays of σ . These power series, when evaluated at certain divisor classes, yield the Todd class of any simplicial toric variety [P3, Theorem 1].

To state the properties of $\mathbf{t}_\sigma$, it will be useful to consider the following variant $\mathfrak{s}$ of the power series $\mathbf{t}$ defined in [P3] by:

$$\mathfrak{s}_\sigma(x_1, \dots, x_n) = \frac{1}{\text{mult}(\sigma)x_1 \cdots x_n} \mathbf{t}_\sigma(x_1, \dots, x_n),$$

which is a Laurent series in $x_1, \dots, x_n$.

The Todd power series $\mathbf{t}_\sigma$ and $\mathfrak{s}_\sigma$ are characterized by the following proposition [P3, Theorem 2], which states that $\mathfrak{s}$ is *additive* under subdivisions (after suitable coordinate changes), and gives the value of $\mathfrak{s}$ on nonsingular cones. An n -dimensional cone is called nonsingular if it is generated by rays forming a basis of the lattice. It is well-known that any cone may be subdivided into nonsingular cones, and that such a subdivision determines a resolution of singularities of the corresponding toric variety (cf. [Fu, Section 2.6].)

Proposition 2.1. *If Γ is a simplicial subdivision of σ then*

$$(11) \quad \mathfrak{s}_\sigma(X) = \sum_{\gamma \in \Gamma(n)} \mathfrak{s}_\gamma(\gamma^{-1}\sigma X).$$

where the sum is taken over the n -dimensional cones of the subdivision Γ . Here X denotes the column vector $(x_1, \dots, x_n)^t$ and we have identified each cone (σ and γ) with the n -by- n matrix whose columns are the coordinates of the rays of that cone.

For nonsingular cones, σ , we have the following expression for $\mathfrak{s}_\sigma$:

$$\mathfrak{s}_\sigma(x_1, \dots, x_n) = \prod_{i=1}^n \frac{1}{1 - e^{-x_i}}.$$

Remark 2.2. It follows immediately that for any cone σ , $\mathfrak{s}_\sigma$ is a rational function of the e^{x_i} .

The Laurent series $\mathfrak{s}_\sigma$ in $x_1, \dots, x_n$ may also be expressed as an exponential sum over the lattice points in the cone, in the spirit of the important earlier work of M. Brion [Br].

Proposition 2.3. *Let $\check{\sigma}$ denote the dual cone in the lattice $M = \text{Hom}(N, \mathbb{Z})$. We then have*

$$\mathfrak{s}_\sigma(x_1, \dots, x_n) = \sum_{m \in \check{\sigma} \cap M} e^{-(\langle m, \rho_1 \rangle x_1 + \dots + \langle m, \rho_n \rangle x_n)},$$

The equality is one of rational functions in the e^{x_i} .

Proof. As noted in [Br], since $\langle m, \rho_i \rangle \geq 0$ for $m \in \check{\sigma} \cap M$, the right hand side has a meaning in the completion of $\mathbb{C}[y_1, \dots, y_n]$ with respect to the ideal $(y_1, \dots, y_n)$, where y_i stands for e^{-x_i} . While it is not obvious, the right hand side is an rational function of the e^{x_i} . See [Br, p.654]. The left hand side is a rational function of the e^{x_i} by Proposition 2.1. By [Br, p.655] and the second formula of Proposition 2.1, these two rational functions are equal on nonsingular cones. As any cone can be subdivided into nonsingular cones, it suffices to verify that the right hand side satisfies the additivity formula of Proposition 2.1. But this follows again from Brion's work. See the proposition of [Br, p.657], for example. Intuitively, this additivity can be seen from the fact that a sum of exponentials over a cone containing a straight line vanishes formally. $\square$

The $\mathfrak{s}_\sigma$ also have an explicit expression in terms of cyclotomic sums, due to Brion and Vergne. Following [BV2], we introduce the following notation. Let $u_1, \dots, u_n$ denote the primitive generators of the dual cone $\check{\sigma}$. Thus we have $\langle u_i, \rho_j \rangle = 0$ if $i \neq j$, and $\langle u_i, \rho_i \rangle \in \mathbb{Z}$, but does not necessarily equal 1. Let N_σ be the subgroup of N generated by the ρ_i , $i = 1, \dots, n$, and let $G_\sigma = N/N_\sigma$. Then G_σ is an abelian group of order $\text{mult}(\sigma)$, which we denote by q . Define characters a_i of G_σ by

$$a_i(g) = e^{2\pi i \frac{\langle u_i, g \rangle}{\langle u_i, \rho_i \rangle}}.$$

Proposition 2.4. *The Laurent series $\mathfrak{s}_\sigma$ coincides with Brion and Vergne's formula expressing their Todd differential operator. Namely, we have*

$$\mathfrak{s}_\sigma(x_1, \dots, x_n) = \frac{1}{q} \sum_{g \in G_\sigma} \prod_{i=1}^n \frac{1}{1 - a_i(g) e^{-x_i}}.$$

Proof. By inspection, the right hand side is a rational function in the variables $y_i = e^{-x_i}$ which takes the value 1 when $y_i = 0$ for all i . Such rational functions embed into the completion of the ring $\mathbb{C}[y_1, \dots, y_n]$ with respect to the ideal $(y_1, \dots, y_n)$. To prove the proposition, it is enough to show that the right hand side and the right hand side of Proposition 2.3 define the same element of this completion. To do so, we expand the right hand side, getting:

$$\frac{1}{q} \sum_{k_1, \dots, k_n \geq 0} \sum_{g \in G_\sigma} \prod_{i=1}^n \left[a_i(g) e^{-x_i} \right]^{k_i},$$

which becomes

$$\frac{1}{q} \sum_{k_1, \dots, k_n \geq 0} e^{-(k_1 x_1 + \dots + k_n x_n)} \sum_{g \in G_\sigma} a_1^{k_1} \dots a_n^{k_n}(g).$$

This last sum over G_σ is either q or 0 , depending on whether $a_1^{k_1} \dots a_n^{k_n}$ is the trivial character of G_σ or not. Comparing the above with the right hand side of Proposition 2, we see that it suffices to show that $a_1^{k_1} \dots a_n^{k_n}$ is the trivial character of G_σ if and only if there exists $m \in \check{\sigma}$ such that $\langle m, \rho_i \rangle = k_i$ for all i . This straightforward lattice calculation is omitted. $\square$

Remark 2.5. The sum in the proposition above appears in the important work of Ricardo Diaz and Sinai Robins [DR]. They use such sums to give an explicit formula for the number of lattice points in a simple polytope. Interestingly, their techniques, which come from Fourier analysis, are seemingly unrelated to the toric geometry discussed above.

2.2. Properties of Todd power series of two-dimensional cones. In this section, we state properties of the power series $\mathbf{t}_\sigma$ for a two-dimensional cone σ .

It is not hard to see that if σ is any two-dimensional cone, then there are relatively prime integers p, q such that σ is lattice-equivalent to the cone

$$\sigma_{(p,q)} \stackrel{\text{def}}{=} \langle (1,0), (p,q) \rangle \subset \mathbb{Z}^2.$$

Here q is determined up to sign and p is determined modulo q . Thus we may arrange to have $q > 0$ and $0 \leq p < q$. This gives a complete classification of two-dimensional cones up to lattice isomorphism. In discussing Todd power series we will abbreviate $\mathbf{t}_{\sigma_{(p,q)}}$ by $\mathbf{t}_{(p,q)}$.

The explicit cyclotomic formula of Proposition 2.4 may be written as:

Proposition 2.6. *The Todd power series of a two-dimensional cone is given by*

$$\mathbf{t}_{(p,q)}(x, y) = \sum_{\omega^q=1} \frac{xy}{(1 - \omega^{-p}e^{-x})(1 - \omega e^{-y})}$$

Proof. With the coordinates above and the notation of Proposition 2.4, we have $u_1 = (q, -p)$, $u_2 = (0, 1)$, and G_σ consists of the lattice points $(0, k)$, $k = 0, \dots, q-1$. The desired equation now follows directly from Proposition 2.4. $\square$

In the two-dimensional case, the additivity formula of Proposition 2.1 can be expressed as an explicit *reciprocity* law. This, and a *periodicity* relation for $\mathfrak{s}$ are contained in the following theorem.

Proposition 2.7. *Let p and q be relatively prime positive integers. Then*

$$(12) \quad \mathfrak{s}_{(p,q)}\left(x - \frac{p}{q}y, \frac{1}{q}y\right) + \mathfrak{s}_{(q,p)}\left(y - \frac{q}{p}x, \frac{1}{p}x\right) = \mathfrak{s}_{(0,1)}(x, y)$$

$$(13) \quad \mathfrak{s}_{(p+q,q)}(x, y) = \mathfrak{s}_{(p,q)}(x, y)$$

Proof. The quadrant $\langle (1,0), (0,1) \rangle$ may be subdivided into cones $\gamma_1 = \langle (1,0), (p,q) \rangle$ and $\gamma_2 = \langle (0,1), (p,q) \rangle$. The cone γ_1 is of type (p,q) , and γ_2 is of type (q,p) . Applying the additivity formula (Proposition 2.1) to this subdivision yields the first equation above.

The second equation follows from the fact that the cones $\langle (1,0), (p,q) \rangle$ and $\langle (1,0), (p+q,q) \rangle$ are lattice isomorphic. $\square$

Let $\mathfrak{s}^{ev}$ (resp. $\mathfrak{s}^{odd}$) denote the part of $\mathfrak{s}$ of even (resp. odd) total degree.

Corollary 2.8. *Given a two-dimensional lattice N , the function $\mathfrak{s} : \mathcal{C}^2(N) \rightarrow \mathbb{Q}((x, y))$ (where $\mathbb{Q}((x, y))$ is the function field of the power series ring $\mathbb{Q}[[x, y]]$) is uniquely determined by properties (12) and (13) and its initial condition $\mathfrak{s}_{(0,1)}(x, y) = 1/((1 - e^{-x})(1 - e^{-y}))$. Since equations (12) and (13) are homogenous with respect to the degrees of x and y , it follows that $\mathfrak{s}^{ev}$ (resp. $\mathfrak{s}^{odd}$) satisfies (12) and (13) with initial condition $\mathfrak{s}_{(0,1)}^{ev}$ (resp. $\mathfrak{s}_{(0,1)}^{odd}$).*

Remark 2.9. Corollary 2.8 has a converse. From equations (12) and (13) it follows that $\mathfrak{s}_{(0,1)}$ satisfies the following relations:

$$\mathfrak{s}_{(0,1)}(x, y) = \mathfrak{s}_{(0,1)}(x - y, y) + \mathfrak{s}_{(0,1)}(y - x, x) \text{ and } \mathfrak{s}_{(0,1)}(x, y) = \mathfrak{s}_{(0,1)}(y, x)$$

Conversely, one can show that given any element $g(x, y) \in \mathbb{Q}((x, y))$ satisfying:

$$g(x, y) = g(x - y, y) + g(y - x, x) \text{ and } g(x, y) = g(y, x)$$

there is a unique function $\mathfrak{g} : \mathcal{C}^2(N) \rightarrow \mathbb{Q}((x, y))$ so that $\mathfrak{g}_{(0,1)} = g$.

2.3. Continued fraction expansion for the Todd power series. In this section, we prove Theorem 3, which expresses the coefficients in the Todd power series of a two-dimensional cone in terms of continued fractions. Before doing so, we first formulate an equivalent version which is more natural from the point of view of toric varieties. The continued fraction expansion in this second version of the formula corresponds directly to a desingularization of the cone.

Given relatively prime integers p, q with $p > 0$ and $0 \leq p < q$, let $a_i, \gamma_i, \delta_i, L_i$ be defined in terms of the negative continued fraction expansions:

$$\frac{q}{q-p} = [a_1, \dots, a_{s-1}], \quad \frac{\gamma_i}{\delta_i} \stackrel{\text{def}}{=} [a_1, \dots, a_{i-1}], \quad L_i \stackrel{\text{def}}{=} \gamma_i x + (q\delta_i + (p-q)\gamma_i)y.$$

We then have the following continued fraction expression for the degree d part $(\mathfrak{t}_\varpi)_d$ of the Todd power series of a two-dimensional cone ϖ .

Theorem 5. *For ϖ a cone of type (p, q) as above, and for $d \geq 2$ an even integer, we have:*

$$(\mathfrak{t}_\varpi)_d(x, y) = -qxy \sum_{i=1}^s P_d(L_{i-1}, L_i) - \lambda_d qxy \sum_{i=1}^{s-1} a_i R_d(L_{i-1}, L_{i+1}) + \lambda_d (xL_1^{d-1} - yL_{s-1}^{d-1})$$

If $d \geq 1$ is odd, then $(\mathfrak{t}_\varpi)_d(x, y) = \frac{1}{2}\lambda_{d-1}q^{d-1}xy(x^{d-2} + y^{d-2})$.

Proof. It will be convenient to choose coordinates so that $\varpi = \langle (0, -1), (q, q-p) \rangle$ (which is easily seen to be lattice equivalent to the cone $\langle (1, 0), (p, q) \rangle$). We now subdivide ϖ into nonsingular cones. It is well-known that for two-dimensional cones this can be done in a canonical way, and that the resulting subdivision has an explicit expression in terms of continued fractions [Fu, Section 2.6]. In our coordinate system, the rays of this nonsingular subdivision of ϖ are given by

$$\begin{aligned} \beta_0 &= (0, -1) \\ \beta_1 &= (1, 0) \\ \beta_2 &= (a_1, 1) \\ &\dots \\ \beta_s &= (q, q-p). \end{aligned}$$

Thus we have $\beta_{i+1} + \beta_{i-1} = a_i \beta_i$, which implies that $\beta_i = (\gamma_i, \delta_i)$.

The cone ϖ is subdivided into cones $\varpi_i = \langle \beta_{i-1}, \beta_i \rangle$, $i = 1, \dots, s$. The N -additivity formula of Proposition 2.7 implies that:

$$\mathfrak{s}_\varpi(x, y) = \sum_{i=1}^s \mathfrak{s}_{\varpi_i}(\varpi_i^{-1} \varpi(x, y)^t),$$

where we again we have identified the 2-dimensional cone ϖ with the 2-by-2 matrix whose columns are the primitive generators of ϖ . One easily sees that this becomes

$$\mathfrak{s}_\varpi(x, y) = \sum_{i=1}^s \mathfrak{s}_{\varpi_i}(-L_{i-1}, L_i)$$

Rewriting the equation in terms of the $\mathfrak{t}_\varpi$ yields

$$\mathfrak{t}_\varpi(x, y) = -qxy \sum_{i=1}^s \frac{\mathfrak{t}_{\varpi_i}(-L_{i-1}, L_i)}{L_{i-1}L_i}$$

Since every ϖ_i is nonsingular, we have

$$\mathfrak{t}_{\varpi_i}(X, Y) = g(X)g(Y)$$

where $g(z) = z/(1 - e^{-z})$. Now consider the degree d part of the above. We will assume $d > 2$ is even and leave the other (easier) cases to the reader. The degree d part of $g(L_{i-1})g(L_i)$ may be written as:

$$-L_{i-1}L_i P_d(-L_{i-1}, L_i) + \lambda_d(L_{i-1}^d + L_i^d).$$

Summing the first term above yields the first term in the equation of the theorem. So it suffices to examine the remaining term:

$$-\lambda_d qxy \sum_{i=1}^s \frac{L_{i-1}^d + L_i^d}{L_{i-1}L_i}.$$

Using the relation $L_{i-1} + L_{i+1} = a_i L_i$, the sum above may be rewritten as

$$\sum_{i=1}^{s-1} a_i \frac{L_{i-1}^{d-1} + L_{i+1}^{d-1}}{L_{i-1} + L_{i+1}} + \left(\frac{L_1^{d-1}}{L_0} + \frac{L_{s-1}^{d-1}}{L_s} \right).$$

Keeping in mind that $L_0 = -qy$ and $L_s = qx$, Theorem 5 follows easily. $\square$

Corollary 2.10. *For a two-dimensional cone $\sigma = \langle \rho_1, \rho_2 \rangle$ of multiplicity q we have:*

$$(14) \quad \mathfrak{t}_\sigma(h_1, h_2) - \frac{1}{2} \mathfrak{t}_{\langle \rho_1 \rangle}(qh_1)h_2 - \frac{1}{2} \mathfrak{t}_{\langle \rho_2 \rangle}(qh_2)h_1 = \mathfrak{t}_\sigma^{ev}(h_1, h_2) - \frac{qh_1 h_2}{2}$$

where $\mathfrak{t}_\sigma^{ev}$ is the even total degree part of the power series $\mathfrak{t}_\sigma$.

Proof. It follows immediately from (and in fact is equivalent to) the formula for the odd part of the Todd power series $\mathfrak{t}_\sigma$ given by Theorem 5. $\square$

We now prove Theorem 3. Let σ be a two-dimensional cone of type (p, q) in a lattice N , and let a_i, h_i, k_i and X_i be as in Theorem 3. The dual cone $\check{\sigma}$ in the dual lattice M is easily seen to have type $(-p, q)$ and so we may choose coordinates in M so that $\check{\sigma} = \langle (0, -1), (q, p) \rangle$ in $\mathbb{Z}^2$. Furthermore, the negative-regular continued fraction expansion of q/p corresponds

naturally to the desingularization of the dual cone $\check{\sigma}$. Explicitly, the desingularization of $\check{\sigma}$ is given by the subdivision

$$\begin{aligned}\rho_0 &= (0, -1) \\ \rho_1 &= (1, 0) \\ \rho_2 &= (b_1, 1) \\ &\dots \\ \rho_r &= (q, p).\end{aligned}$$

One has $\rho_{i+1} + \rho_{i-1} = b_i \rho_i$ and thus $\rho_i = (h_i, k_i)$. Applying Theorem 5 to $\check{\sigma}$ expresses $\mathbf{t}_{(-p,q)}$ in terms of the b_i . However $\mathbf{t}_{(-p,q)}$ is related to $\mathbf{t}_{(p,q)}$ via the relation

$$\mathbf{t}_{(p,q)}(x, y) = \mathbf{t}_{(-p,q)}(-x, y) + \frac{qxy}{1 - e^{-qy}}.$$

In this way, we obtain an expression for $\mathbf{t}_{(p,q)}$ in terms of the b_i , which concludes the proof of Theorem 3.

2.4. Zeta function values in terms of Todd power series. In this section, we prove Theorem 2 which expresses values of the zeta function of a two-dimensional cone in terms of the Todd power series. Three ingredients are involved in the proof of this theorem: an asymptotic series formula (Proposition 2.11), a polytope summation formula (Proposition 2.12) and a cone summation formula (Proposition 2.13). We begin with the first ingredient:

Lemma 2.11. [Za4, Proposition 2] *Let $\phi(s) = \sum_{\lambda>0} a_\lambda \lambda^{-s}$ be a Dirichlet series where $\{\lambda\}$ is a sequence of positive real numbers converging to infinity. Let $E(t) = \sum_{\lambda>0} a_\lambda e^{-\lambda t}$ be the corresponding exponential series. Assume that $E(t)$ has the following asymptotic expansion as $t \rightarrow 0$:*

$$(15) \quad E(t) \sim \sum_{n=-1}^{\infty} c_n t^n$$

Then it follows that

- $\phi(s)$ can be extended to a meromorphic function on $\mathbb{C}$.
- $\phi(s)$ has a simple pole at $s = 1$, and no other poles.
- The values of ϕ at nonpositive integers are given by: $\phi(-n) = (-1)^n n! c_n$.

We now present our second ingredient, a polytope summation formula. This proposition is a variant of the lattice point formula of [BV2]. A polytope of dimension n is called *simple* if each of its vertices lies on exactly n facets ($n - 1$ -dimensional faces) of the polytope.

Proposition 2.12. *Let N be an n -dimensional lattice, P a simple lattice polytope in M and Σ its associated fan in N . For every analytic function $\phi : M_{\mathbb{R}} \rightarrow \mathbb{R}$, we have the following asymptotic expansion as $t \rightarrow 0$:*

$$(16) \quad \sum_{a \in P \cap M} \phi(ta) \sim \mathbf{t}_{\Sigma} \left(\frac{\partial}{\partial h} \right) \diamond \int_{P(h)} \phi(tu) du$$

where $\mathbf{t}_{\Sigma}$ is the Todd power series of [BV2, Definition 10].

Proof. First of all, the meaning of the right hand side is as follows: we consider the degree k Taylor expansion $\phi = \phi_k + R_k$ of ϕ , where ϕ_k is a polynomial in $M_{\mathbb{R}}$ of degree k and R_k is the remainder satisfying $\lim_{a \rightarrow 0} |a|^k R_k(a) = 0$. It follows that $\int_{P(h)} P_k(tu) du$ is a polynomial in t and h of degree k (with respect to t) and that $\int_{P(h)} R_k(tu) du = o(t^k)$ at $t = 0$ (with the notation that $f(t) = o(t^k)$ if and only if $\lim_{t \rightarrow 0} f(t)t^{-k} = 0$). Thus,

$$\mathbf{t}_{\Sigma} \left(\frac{\partial}{\partial h} \right) \diamond \int_{P(h)} \phi(tu) du = \mathbf{t}_{\Sigma} \left(\frac{\partial}{\partial h} \right) \diamond \int_{P(h)} \phi_k(tu) du + o(t^k).$$

On the other hand,

$$\begin{aligned} \sum_{a \in P \cap M} \phi(ta) &= \sum_{a \in P \cap M} \phi_k(ta) + \sum_{a \in P \cap M} R_k(ta) \\ &= \sum_{a \in P \cap M} \phi_k(ta) + o(t^k). \end{aligned}$$

Brion-Vergne [BV2, theorem 11] prove that for every polynomial function (such as ϕ_k) on $M_{\mathbb{R}}$ we have:

$$\sum_{a \in P \cap M} \phi_k(ta) = \mathbf{t}_{\Sigma} \left(\frac{\partial}{\partial h} \right) \diamond \int_{P(h)} \phi_k(tu) du,$$

which concludes the proof. $\square$

We call a function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ *rapidly decreasing* if it is analytic and for every constant coefficients differential operator D , and every subset I of $[n] = \{1, \dots, n\}$, the restriction $D(\phi)|_I$ obtained by setting $x_i = 0$ for $i \notin I$ is in $L^1(\mathbb{R}_+^I)$. Examples of rapidly decreasing functions can be obtained by setting $\phi = P \exp(Q)$ where P is a polynomial on $\mathbb{R}^n$ and $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ is totally positive, i.e., its restriction to $\mathbb{R}_+^I$ takes positive values for every subset I of $[n]$.

Proposition 2.13. *Let N be an n -dimensional lattice. For every $\sigma \in \mathcal{C}^n(N)$, and every rapidly decreasing analytic function $\phi : M_{\mathbb{R}} \rightarrow \mathbb{R}$, we have the following asymptotic expansion as $t \rightarrow 0$:*

$$(17) \quad \sum_{a \in \tilde{\sigma} \cap M} \phi(ta) \sim \mathbf{t}_{\sigma} \left(\frac{\partial}{\partial h} \right) \diamond \int_{\tilde{\sigma}(h)} \phi(tu) du$$

Proof. First of all, the right hand side of the above equation has the following meaning: consider the decomposition $\mathbf{t}_{\sigma} = \sum_k \mathbf{t}_{\sigma,k}$ of the power series $\mathbf{t}_{\sigma}$, where $\mathbf{t}_{\sigma,k}$ is a homogenous polynomial of degree k . A change of variables $v = tu$ implies that

$$\mathbf{t}_{\sigma,k} \left(\frac{\partial}{\partial h} \right) \diamond \int_{\tilde{\sigma}(h)} \phi(tu) du = \mathbf{t}_{\sigma,k} \left(\frac{\partial}{\partial h} \right) \diamond \int_{\tilde{\sigma}(th)} \phi(v) dv / t^n = t^{k-n} \mathbf{t}_{\sigma,k} \left(\frac{\partial}{\partial h} \right) \diamond \int_{\tilde{\sigma}(h)} \phi(v) dv$$

is a multiple of t^{k-n} , thus the right hand side is defined to be the Laurent power series in t given by

$$\sum_{k=0}^{\infty} t^{k-n} \mathbf{t}_{\sigma,k} \left(\frac{\partial}{\partial h} \right) \diamond \int_{\tilde{\sigma}(h)} \phi(v) dv.$$

For the proof of the proposition, truncate in some way the cone $\check{\sigma}$ in M to obtain a simple convex polytope P , with associated fan Σ . Since $\cup_{r>0} rP = \check{\sigma}$, using the convergence properties of ϕ , we obtain as $r \rightarrow \infty$:

$$\begin{aligned}
\sum_{a \in \check{\sigma} \cap M} \phi(ta) &= \lim_r \sum_{a \in rP \cap M} \phi(ta) \\
&\sim \lim_r \mathbf{t}_\Sigma \left(\frac{\partial}{\partial h} \right) \diamond \int_{(rP)(h)} \phi(tu) du \\
&= \lim_r \mathbf{t}_\Sigma \left(\frac{\partial}{\partial h} \right) \diamond \int_{(rtP)(th)} \phi(v) dv / t^n \\
&= \lim_r \sum_k \mathbf{t}_{\Sigma,k} t^{k-n} \left(\frac{\partial}{\partial h} \right) \diamond \int_{(rtP)(h)} \phi(v) dv \\
&= \sum_k \lim_r \mathbf{t}_{\Sigma,k} t^{k-n} \left(\frac{\partial}{\partial h} \right) \diamond \int_{(rtP)(h)} \phi(v) dv \\
&= \sum_k \mathbf{t}_{\sigma,k} t^{k-n} \left(\frac{\partial}{\partial h} \right) \diamond \int_{P(h)} \phi(v) dv,
\end{aligned}$$

which concludes the proof. $\square$

In case of a two-dimensional lattice N and a rapidly decreasing function $\phi : M_{\mathbb{R}} \rightarrow \mathbb{R}$, using the weight function wt of equation (2) and inclusion-exclusion, we obtain the following

Corollary 2.14. *For a two-dimensional cone $\sigma = \langle \rho_1, \rho_2 \rangle$ of multiplicity q in N , we have the asymptotic expansion as $t \rightarrow 0$:*

$$(18) \quad \sum_{a \in \check{\sigma} \cap M} wt(\check{\sigma}, a) \phi(ta) \sim \left\{ \mathbf{t}_\sigma^{ev} \left(\frac{\partial}{\partial h_1}, \frac{\partial}{\partial h_2} \right) - \frac{q}{2} \frac{\partial^2}{\partial h_1 \partial h_2} \right\} \diamond \int_{\check{\sigma}(h)} \phi(tu) du,$$

where the right hand side lies in the formal power series ring $t^{-2} \mathbb{R}[[t^2]]$.

Proof of Theorem 2. Recall that τ is a cone in M , σ is its dual in N and Q is homogenous quadratic, totally positive on τ ; thus e^{-Q} is rapidly decreasing on $M_{\mathbb{R}}$. Corollary 2.14 implies that the generating function

$$(19) \quad Z_{Q,\tau}(t) = \sum_{a \in \tau \cap M} wt(\tau, a) e^{-tQ(a)} = \sum_{a \in \tau \cap M} wt(\tau, a) e^{-Q(t^{1/2}a)},$$

satisfies the hypothesis of Lemma 2.11, which in turn yields Theorem 2. $\square$

Remark 2.15. Notice that the above proof of Theorem 2 used crucially the fact that $B_1 = -\frac{1}{2}$ and the definition of the weight function wt . If we had weighted the sum defining the zeta function in any other way, the resulting variation of Theorem 2 would not hold.

3. ZETA FUNCTIONS OF NUMBER FIELDS

3.1. Parametrizing triples (M_n, Q_b, τ_b) . In this section, we review in detail the construction of the triple (M_b, Q_b, τ_b) mentioned in Section 1.1 that is associated to a sequence $b = (b_0, \dots, b_{r-1})$ of integers greater than 1 and not all equal to 2. Our notation and construction is borrowed from [Za4]. Given a sequence b as above, we extend it to a sequence of

integers parametrized by the integers by defining $b_k = b_{k \bmod r}$. Furthermore, for an integer k , we define

$$w_k = \llbracket b_k, \dots, b_{k+r-1} \rrbracket = b_k - \frac{1}{b_{k+1} - \frac{1}{b_{k+2} - \dots}}$$

where $\llbracket b_k, \dots, b_{k+r-1} \rrbracket$ denotes the infinite continued fraction with period r . Note that $w_k = w_{k+r}$ for all integers k , and that, by definition,

$$w_0 = b_0 - \frac{1}{b_1 - \frac{1}{\dots b_{r-1} - \frac{1}{w_0}}}$$

from which it follows that w_0 satisfies a quadratic equation $A_b w^2 + B_b w + C_b = 0$ where A_b, B_b, C_b are (for a fixed r) polynomials in b_i with integer coefficients.² Let $D_b = B_b^2 - 4A_b C_b$ be the discriminant. Since $b_i > 1$, it follows that $D_b > 0$, and that the roots of the quadratic equation are $w_0 = (-B_b + \sqrt{D_b})/(2A_b)$ and $w'_0 = (-B_b - \sqrt{D_b})/(2A_b)$. We thus have that $w_0 > 1 > w'_0$. We define the complex numbers $A_0 = 1$, $A_{k-1} = A_k w_k$ for $k \in \mathbb{Z}$. Since $w_k = b_k - \frac{1}{w_{k+1}}$ it follows that

$$(20) \quad A_{k-1} + A_{k+1} = b_k A_k$$

for all integers k . Let us define $M_b = \mathbb{Z}w_0 + \mathbb{Z}$ to be the rank two lattice with the oriented basis $\{w_0, 1\}$. Note that M_b is a rank two lattice in the real quadratic field $K_b = \mathbb{Q}(\sqrt{D_b})$. Due to the fact that $A_{-1} = w_0$ and the recursion relation (20), it follows that $M_b = \mathbb{Z}A_k + \mathbb{Z}A_{k+1}$ for all integers k , and thus $\epsilon M_b = M_b$ where $\epsilon = w_0 \dots w_{r-1}$. Thus $U_b = \{\epsilon^m | m \in \mathbb{Z}\}$ acts on M_b by multiplication, and the zeta function of (the inverse of) the narrow ideal class of M_b satisfies

$$\zeta(M_b, s) = N(M_b)^s \sum_{\substack{a \in M_b/U_b \\ a \gg 0}} \frac{1}{Q_b(a)^s},$$

where in the above summation we exclude 0, and $N(M_b)$ is a constant, defined in [Za4], which is not so crucial for our purposes. We are therefore interested to describe the quotient M_b/U_b . The action of U_b on M_b implies that $\epsilon A_k = A_{k-r}$. The matrix of multiplication by ϵ on M_b , in terms of the basis $\{\beta_1, \beta_2\} \stackrel{\text{def}}{=} \{A_{-1}, A_0\}$ can be calculated as follows. Recall first that in terms of the above basis of M_b , we have that:

$$(21) \quad A_k = -p_k A_{-1} + q_k A_0 \text{ where } \frac{p_k}{q_k} = [b_0, \dots, b_{k-1}]$$

which implies that

$$\epsilon^{-1} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} A_{r-1} \\ A_r \end{pmatrix} = \begin{pmatrix} -p_{r-1} & -p_r \\ q_{r-1} & q_r \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix},$$

from which it follows that the matrix of ϵ is given by

$$(22) \quad \begin{pmatrix} q_r & p_r \\ -q_{r-1} & -p_{r-1} \end{pmatrix} = \begin{pmatrix} b_0 q - p & q \\ -b_0 p' + (pp' - 1)/q & -p' \end{pmatrix} = \begin{pmatrix} b_0 & -1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} b_{r-1} & -1 \\ 1 & 0 \end{pmatrix}$$

²not to be confused with $A_0, \dots, A_r$ introduced below, despite the rather poor choice of notation.

where

$$(23) \quad \frac{q}{p} \stackrel{\text{def}}{=} [b_1, \dots, b_{r-1}],$$

(with the understanding that $q = 1, p = 0$ if $r = 1$), $p' = \text{numerator}[b_1, \dots, b_{r-2}]$, which is a multiplicative inverse of $p \bmod q$ (with the understanding that $p' = 0$ (resp. 1) if $r = 1$ (resp. 2)).

Lemma 3.1. *The orbit space M_b/U_b has as a fundamental domain a semiopen two-dimensional simplicial cone whose closure is the cone $\langle A_0, A_r \rangle$ in K_b . Furthermore, the (closed) cone $\tau_b = \langle A_0, A_r \rangle$ in M_b is of type $(-p, q)$ and the dual cone σ_b in N_b is of type (p, q) where $0 \leq p < q$ as in equation (23) above.*

Proof. From Figure 1, it follows that a fundamental domain of M_b/U_b is a semiopen two-dimensional cone whose closure is the cone $\tau_b = \langle A_r, A_0 \rangle$ in M_b . The cone $\langle A_r, A_0 \rangle$ is canonically subdivided into nonsingular cones $\langle A_{i+1}, A_i \rangle$ (for $i = 0, \dots, r-1$), and using equation (20), it follows that τ_b is of type (c_1, q) where $0 \leq c_1 < q$ and $\frac{q}{q-c_1} = [b_1, \dots, b_{r-1}]$. Therefore the dual cone σ_b in N is of type (p, q) where $p = q - c_1$. Thus (p, q) satisfies equation (23). $\square$




Figure 1. A fundamental domain of M_b/U_b .

Since $M_b \subseteq \mathbb{Q}(\sqrt{D_b})$ is a rank two lattice of a real quadratic field, we can define a homogenous function $Q_b : M_{\mathbb{R}} \rightarrow \mathbb{R}$ by

$$Q_b(xw_0 + y) = C_b x^2 - B_b xy + A_b y^2$$

This agrees with the (normalized) norm function of the real quadratic field $\mathbb{Q}(\sqrt{D_b})$ considered by Zagier [Za4, p 138]. We close this section with the following:

Proposition 3.2. *Let $l, m \in \mathbb{Z}$ with $l \neq m$. Then the coefficients of x^2, xy and y^2 of the quadratic form $Q_b(xA_l + yA_m)$ are given by polynomials in b_i with integer coefficients.*

Proof. We first claim that in terms of the basis $\{A_r, A_0\}$ we have:

$$(24) \quad Q_b(xA_r + yA_0) = qQ_{\Theta}(x, y)$$

where $Q_{\Theta}(x, y) = x^2 + \Theta xy + y^2$ and $\Theta = b_0 q - p - p'$.

The above claim follows from the fact that A_b, B_b, C_b can be calculated (using their definition) in terms of the b_i as follows:

$$A_b = q, \quad B_b = -b_0 q + p - p', \quad \text{and} \quad C_b = q_{r-1} = b_0 p' + (1 - pp')/q,$$

together with a change of variables formula from $\{A_{-1}, A_0\}$ to $\{A_r, A_0\}$ given by equation (21).

We then claim that if l, m are distinct integers, then

$$(25) \quad Q_b(xA_l + yA_m) = \frac{1}{q} Q_\Theta(H_l x + H_m y, K_l x + K_m y)$$

where $\frac{h_i}{k_i} = [b_1, \dots, b_{i-1}]$, $H_i x + K_i y \stackrel{\text{def}}{=} X_i \stackrel{\text{def}}{=} -h_i x + (qk_i - ph_i)y$.

This follows from equation (24) using the change of variables given in equation (21).

We now claim that the coefficients of x^2, xy and y^2 in equation (25) are polynomials in the b_i with integer coefficients.

Indeed, observe that h_i, k_i are polynomials in b_i with integer coefficients. Since $H_i = -h_i, K_i = qk_i - ph_i = -ph_i \pmod{q}$, it follows that:

$$\begin{aligned} Q_\Theta(H_l x + H_m y, K_l x + K_m y) &= (h_l x + h_m y)^2 + \Theta(h_l x + h_m y)(ph_l x + ph_m y) \\ &\quad + (ph_l x + ph_m y)^2 \pmod{q} \\ &= (h_l x + h_m y)^2 (1 + \Theta p + p^2) \pmod{q}. \end{aligned}$$

Since $\Theta = qb_0 - p - p'$, and $pp' \equiv 1 \pmod{q}$, it follows that $1 + \Theta p + p^2 = 0 \pmod{q}$, which finishes the proof of the proposition. $\square$

Remark 3.3. The notation of Theorem 3 (and its proof, in Section 2.3) corresponds to the notation of this section. Indeed, the cone σ_b in N is of type (p, q) where p, q are given in terms of equation (23). The rays ρ_i (for $i = 1, \dots, r-1$) that desingularize the cone τ_b in M are given by $\rho_i = A_i$. In addition, the h_i, k_i, X_i that appear in Theorem 3 match the notation Proposition 3.2.

3.2. Proof of Theorem 1.

Proof. The main idea is to use Theorem 2 which calculates the zeta values in terms of the Todd operator of σ_b , and Theorem 3 which expresses the Todd operator in terms of the b_i . The expression that we obtain for the zeta values differ from the one of equation (4) by an error term, which vanishes identically, as one can show by an explicit calculation.

Now, for the details, we follow the notation of Section 3.1. We begin by calculating the integral $\int_{\tau_b(h_1, h_2)} e^{-Q(u)} du$.

Using the parametrization $\mathbb{R}^2 \rightarrow M_{\mathbb{R}}$ given by: $(w_1, w_2) \rightarrow w_1 A_r + w_2 A_0$, it follows that the preimage of $\tau_b(x, y)$ in $\mathbb{R}^2$ is given by $\{(w_1, w_2) | w_1 \geq -x/q, w_2 \geq -y/q\}$. Thus we have:

$$\int_{\tau_b(x, y)} e^{-Q_b(u)} du = q \int_{w_1 = -x/q}^{\infty} \int_{w_2 = -y/q}^{\infty} e^{-Q_b(qw_1, -pw_1 + w_2)} dw_2 dw_1$$

Differentiating, we get

$$\left(\frac{\partial}{\partial x} \right) \left(\frac{\partial}{\partial y} \right) \diamond \int_{\tau_b(x, y)} e^{-Q_b(u)} du = \frac{1}{q} e^{-Q_b(-xA_r/q - yA_0/q)} = \frac{1}{q} e^{-1/q Q_\Theta(x, y)}$$

Using the following elementary identity:

$$\left(\alpha \frac{\partial}{\partial \bar{x}} + \beta \frac{\partial}{\partial \bar{y}} \right)^i \left(\gamma \frac{\partial}{\partial \bar{x}} + \delta \frac{\partial}{\partial \bar{y}} \right)^j \Big|_{\substack{\bar{x} = \alpha a + \gamma b \\ \bar{y} = \beta a + \delta b}} f(\bar{x}, \bar{y}) = \left(\frac{\partial}{\partial x} \right)^i \left(\frac{\partial}{\partial y} \right)^j \Big|_{\substack{x = a \\ y = b}} f(\alpha x + \gamma y, \beta x + \delta y)$$

(and temporarily abbreviating $\frac{\partial}{\partial x}$ by x) we obtain that

$$\begin{aligned}
(qxyX_l^aX_m^b) \diamond \int_{\tau_b(x,y)} e^{-Q_b(u)} du &= X_l^aX_m^b \diamond e^{-1/qQ_\Theta(x,y)} \\
&= \left(\frac{\partial}{\partial x}\right)^a \left(\frac{\partial}{\partial y}\right)^b \diamond e^{-1/qQ_\Theta(H_lx+H_my,K_lx+K_my)} \\
&= \left(\frac{\partial}{\partial x}\right)^a \left(\frac{\partial}{\partial y}\right)^b \diamond e^{-Q_b(xA_l+yA_m)}, \quad \text{as well as}
\end{aligned}$$

$$\begin{aligned}
(xX_1^{2n+1} + yX_{r-1}^{2n+1}) \diamond \int_{\tau_b(x,y)} e^{-Q_b(u)} du &= \\
\frac{1}{q^{n+1}} \int_0^\infty \left(\left(\frac{\partial}{\partial x} + p\frac{\partial}{\partial y}\right)^{2n+1} + \left(\frac{\partial}{\partial x} + p'\frac{\partial}{\partial y}\right)^{2n+1} \right) \Big|_{x=0} e^{-Q_\Theta(x,y)} dy
\end{aligned}$$

The above, together with Theorem 3, implies that:

$$\begin{aligned}
\zeta_{Q_b, \tau_b}(-n) &= (-1)^n n! \left\{ \sum_{i=1}^r P_{2n+2} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) \diamond e^{-Q_b(xA_{i-1}+yA_i)} \right. \\
&\quad \left. + \lambda_{2n+2} \sum_{i=1}^r b_i R_{2n+2} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) \diamond e^{-Q_b(xA_{i-1}+yA_{i+1})} + E_{2n+2}(b) \right\}
\end{aligned}$$

where

$$\begin{aligned}
E_{2n+2}(b) &= \lambda_{2n+2} \left\{ -b_0 R_{2n+2} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) e^{-1/q(Lx^2+Mxy+Ny^2)} \right. \\
&\quad \left. - \frac{1}{q^{n+1}} \int_0^\infty \left(\left(\frac{\partial}{\partial x} + p\frac{\partial}{\partial y}\right)^{2n+1} + \left(\frac{\partial}{\partial x} + p'\frac{\partial}{\partial y}\right)^{2n+1} \right) \Big|_{x=0} e^{-Q_\Theta(x,y)} dx \right\}
\end{aligned}$$

and

$$L = qp'b_0 + 1 - pp', \quad M = qb_0\Theta_b + 2(pp' - 1), \quad N = qpb_0 + 1 - pp'.$$

Fixing r , the length of the sequence $(b_0, \dots, b_{r-1})$, Lemma 3.4 below can be used to express $E_{2n+2}(b)$ as a polynomial in $1/q, p, p', b_0$. An explicit but lengthy calculation implies that $E_{2n+2}(b) = 0$ for any b . We have thus succeeded in expressing $\zeta_{Q_b, \sigma_b}(-n)$ as a polynomial in b_i with rational coefficients, symmetric under cyclic permutation, (since $Q_b(xA_l + yA_m)$ is symmetric under cyclic permutation). This concludes the proof of Theorem 1. Equation (6) follows easily using $\lambda_1 = 1/2, \lambda_2 = 1/12$. $\square$

Lemma 3.4. [Za4] *For every $i, j \geq 0$ with $i + j$ even, we have:*

$$\left(\frac{\partial}{\partial x}\right)^i \left(\frac{\partial}{\partial y}\right)^j \diamond e^{-(ax^2+bxy+cy^2)} = i!j!(-1)^{(i+j)/2} a^{i/2} c^{j/2} \sum_{i_2=0; i_2 \equiv i \pmod{2}}^{\min\{i,j\}} \frac{(a^{-1/2}bc^{-1/2})^{i_2}}{((i-i_2)/2)!i_2!((j-i_2)/2)!}$$

Furthermore, for every $n \geq 1$ we have:

$$\int_{y=0}^{\infty} \left(\frac{\partial}{\partial x} \right)^{2n-1} \diamond e^{-(ax^2+bx+cy^2)} dy = -\frac{(2n-1)!}{2c^n} \sum_{r=0}^{n-1} (-1)^r \frac{(n-1-r)!}{r!(2n-1-2r)!} a^r b^{2n-1-2r} c^r$$

4. DEDEKIND SUMS IN TERMS OF TODD POWER SERIES

In this section we give a proof of Theorem 4. To do this we will use Proposition 2.3, which can be used to express the coefficients $f_{i,j}$ as a sum of rational numbers, which turn out to equal products of certain values of the Benoulli polynomials. (Note, in contrast, that Proposition 2.4 expresses this same number in terms of roots of unity instead of rational numbers.)

Let p and q be as in the statement of Theorem 4, and let σ be the cone $\langle (1,0), (p,q) \rangle$ in $\mathbb{Z}^2$. We let $\rho_1 = (1,0)$, and $\rho_2 = (p,q)$ denote the generators of this cone. The left hand side of Theorem 4, $f_{i,j}(p,q)$, equals the coefficient of $x^i y^j$ in the power series $\mathbf{t}_\sigma(x,y)$. We now compute this power series using Proposition 2.3. This proposition contains an expansion for the power series $\mathbf{s}_\sigma$, which is equivalent to the following expansion of $\mathbf{t}_\sigma$:

$$\mathbf{t}_\sigma(x,y) = qxy \sum_{m \in \check{\sigma} \cap M} e^{-(\langle m, \rho_1 \rangle x + \langle m, \rho_2 \rangle y)}.$$

Every point of $\check{\sigma} \cap M$ can be written uniquely as a nonnegative integral combination of the generators $u_1 = (q, -p)$ and $u_2 = (0, 1)$ of $\check{\sigma}$, plus a lattice point in the semiopen parallelepiped

$$P = \{cu_1 + du_2 | c, d \in [0, 1)\}.$$

Using $\langle u_1, \rho_1 \rangle = \langle u_2, \rho_2 \rangle = q$, it follows that

$$\mathbf{t}_\sigma(x,y) = qxy \frac{1}{1 - e^{-qx}} \frac{1}{1 - e^{-qy}} \sum_{m \in P \cap M} e^{-(\langle m, \rho_1 \rangle x + \langle m, \rho_2 \rangle y)}.$$

One finds also that

$$P = \left\{ \left(k, \left\{ \frac{pk}{q} \right\} - \frac{pk}{q} \right) : k = 0, \dots, q-1 \right\},$$

where $\{x\} \in [0, 1)$ denotes the fractional part of x . (Note this is slightly different from $\langle x \rangle \in (0, 1]$, which appears in the definition of $s_{i,j}$: by definition, $\langle 0 \rangle = 1$, while $\{0\} = 0$.) One then finds that

$$\mathbf{t}_\sigma(x,y) = qxy \frac{1}{1 - e^{-qx}} \frac{1}{1 - e^{-qy}} \sum_{k=0}^{q-1} e^{-q(\frac{k}{q}x + \{\frac{kp}{q}\}y)}.$$

We may then compute $f_{i,j}(p,q)$ as the coefficient of $x^i y^j$ in the above expression. It is convenient to replace x and y with $-x$ and $-y$, which introduces a factor of $(-1)^{i+j}$. We obtain

$$f_{i,j}(p,q) = (-1)^{i+j} q^{i+j-1} \sum_{k=0}^{q-1} \text{coeff} \left(x^i; \frac{x e^{\frac{k}{q}x}}{1 - e^x} \right) \text{coeff} \left(y^j; \frac{y e^{\{\frac{kp}{q}\}y}}{1 - e^y} \right).$$

It is then clear that we can write the above sum in terms of values of the Bernoulli polynomials, as follows:

$$f_{i,j}(p, q) = (-1)^{i+j} q^{i+j-1} \sum_{k=0}^{q-1} \text{Ber}_i\left(\frac{k}{q}\right) \text{Ber}_j\left(\left\{\frac{kp}{q}\right\}\right).$$

Using the identity

$$\text{Ber}_j(\lambda) = (-1)^j \text{Ber}_j(1 - \lambda),$$

we may rewrite our expression as

$$f_{i,j}(p, q) = (-1)^i q^{i+j-1} \sum_{k=0}^{q-1} \text{Ber}_i\left(\frac{k}{q}\right) \text{Ber}_j\left(\left\langle -\frac{kp}{q} \right\rangle\right).$$

The sum on the right hand side is easily seen to equal the sum defining $s_{i,j}(p, q)$, except for a possible discrepancy in the $k = 0$ term. It follows easily from the definitions that the $k = 0$ terms actually match unless $i = j = 1$, or $i = 1$ and j is even, or $j = 1$ and i is even. In these cases, we need the correction terms which appear in the statement of the Theorem.

REFERENCES

- [AV] T. Apostol, T. Vu, *Identities for sums of Dedekind type*, J. Number Theory **14** (1982) 391–396.
- [Ba] A. I. Barvinok, *A polynomial time algorithm for counting integral points in polyhedra when the dimension is fixed*, Math. Oper. Res. **19** (1994) no.4 769–779.
- [Br] M. Brion, *Points entier dans les polyèdres convexes*, Ann. Scient. Éc. Norm. Sup. **21** (1988) no.4 653–663.
- [BV1] M. Brion and M. Vergne, *An equivariant Riemann-Roch theorem for simplicial toric varieties*, J. Reine Angew. Math **482** (1997) 67–92.
- [BV2] ———, *Lattice points in simple polytopes*, Journal AMS **10** (1997) 371–392.
- [CS] S. Cappell, J. Shaneson, *Genera of algebraic varieties and counting lattice points*, Bull. Amer. Math. Soc. **30** (1994), 62–69.
- [C-N1] P. Cassou-Noguès, *Valeurs aux entiers négatifs des fonctions zêta et fonctions zêta p-adiques*, Inventiones Math. **51** (1979) 29–59.
- [C-N2] ———, *Valeurs aux entiers négatifs des séries de Dirichlet associées a un polynôme I*, Journal of Number Theory **14** (1982) 32–64.
- [C-N3] ———, *Valeurs aux entiers négatifs des séries de Dirichlet associées a un polynôme II, III*, Amer. J. Math **106** (1984) 255–299 and **109** (1987) 71–89.
- [Da] V. I. Danilov, *The geometry of toric varieties*, Russian Math. Surveys **33** (1978), 97–154.
- [DR] R. Diaz, S. Robins *The Ehrhart polynomial of a lattice polytope*, Annals of Math. **145** (1997) 503–518.
- [Fu] W. Fulton, *Introduction to toric varieties*, Annals of Math. Studies **131** Princeton, New Jersey 1993.
- [Gu] V. Guillemin, *Riemann-Roch for toric orbifolds*, J. Diff. Geom. **45** (1997) 53–73.
- [Ha] U. Halbritter, *Some new reciprocity formulas for generalized Dedekind sums*, Results Math. **8** (1985) 21–46.
- [Hs] D. Hayes, *Brumer elements over a real quadratic base field*, Exposition. Math. **8** no.2 (1990) 137–184.
- [KK] J.M. Kantor, A.G. Khovanskii *Integral points in convex polyhedra, combinatorial Riemann-Roch and generalized MacLaurin formulae*, IHES (1992) 932–937.
- [KP] A.G. Khovanskii, A.V. Pukhlikov, *A Riemann-Roch theorem for integrals and sums of quasipolynomials over virtual polytopes*, St. Petersburg Math. J. **4** (1993) 789–912.
- [Kl] H. Klingen, *Über die Werke der Dedekindsche Zeta Funktion*, Math. Ann. **145** (1962) 265–272.
- [Li] S. Lichtenbaum, *Talk at I.A.S.*, mimeographed notes 1971.

- [Me] C. Meyer, *Über die Berechnung der Klassenzahl abelscher Körper quadratischen Zahlkörpern*, Akademie-Verlag Berlin 1955.
- [Mo] R. Morelli, *Pick's Theorem and the Todd class of a toric variety*, Adv. Math. **100:2** (1993), 183-231.
- [P1] J. E. Pommersheim, *Toric varieties, lattice points and Dedekind sums*, Math. Ann. **295** (1993) 1-24.
- [P2] ———, *Products of cycles and the Todd class of a toric variety*, Journal of AMS **9** (1996) no. 3, 813-826.
- [P3] ———, *Barvinok's algorithm and the Todd class of a toric variety*, J. Pure Appl. Alg., **117 & 118** (1997), 519-533.
- [R-G] H. Rademacher, E. Grosswald, *Dedekind sums*, Carus Math. Monographs, **16** Mathematical Association of America, Washington D.C., 1972.
- [Sc1] R. Sczech, *Eisenstein cocycles for $GL_2(\mathbb{Q})$ and values of L -functions in real quadratic fields*, Comment. Math. Helvetici, **67** (1992) 363-382.
- [Sc2] ———, *Eisenstein group cocycles for $GL_n(\mathbb{Q})$ and values of L -functions*, Inventiones **113** (1993) 581-616.
- [Sh1] T. Shintani, *On evaluation of zeta functions of totally real algebraic number fields at nonpositive integers*, J. Fac. Sci. Univ. Tokyo, Sec. 1A, **23** (1976) 393-417.
- [Sh2] ———, *On special values of zeta functions of totally real algebraic number fields*, Proc. Intern. Congress of Mathematicians, Helsinki (1978) 591-597.
- [Si1] C. L. Siegel, *Bernoulische Polynome und Quadratische Zahlkörper*, Nachr. Akad. Wiss. Göttingen Math. Physik. **3** (1970) 7-38.
- [Si2] ———, *Über die Fourrierschen Koeffizienten von Modulformen*, Nachr. Akad. Wiss. Göttingen Math. Physik. **3** (1970) 15-56.
- [So] D. Solomon, *Algebraic properties of Shintani's generating functions: Dedekind sums and cocycles on $PGL_2(\mathbb{Q})$* , preprint July 1995.
- [St] G. Stevens, *The Eisenstein measure and real quadratic fields*, in the proceedings of the International Number Theory Conference (Université Laval, 1987) J.M. de Koninck and C. Levesque eds., de Gruyter, 1989.
- [Za1] D. Zagier, *Higher dimensional Dedekind sums*, Math. Ann. **202** (1973) 149-172.
- [Za2] ———, *A Kronecker limit formula for real quadratic fields*, Math. Ann. **213** (1975) 153-182.
- [Za3] ———, *Nombres de classes et fractions continues*, Soc. Math. France Asterisque, **24-25** (1975) 81-97.
- [Za4] ———, *Valeurs des fonctions zêta des corps quadratiques réels aux entiers négatifs*, Soc. Math. France Asterisque, **41-42** (1977) 135-151.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA.

E-mail address: stavros@math.harvard.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NM 88003, USA.

E-mail address: jamie@math.nmsu.edu